#### § 520.48

#### §520.48 Altrenogest solution.

- (a) Specifications. Each milliliter (mL) of solution contains 2.2 milligrams (mg) altrenogest.
- (b) Sponsor. See No. 057926 in  $\S510.600$ (c) of this chapter.
- (c) Tolerances. See §556.36 of this chapter.
- (d) Conditions of use—(1)*Horses*—(i)*Amount*. 1.0 mL per 110 pounds body weight (0.044 mg/kg) daily for 15 consecutive days.
- (ii) *Indications for use*. For suppression of estrus in mares.
- (iii) *Limitations*. For oral use in horses only; avoid contact with the skin. Do not administer to horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Swine—(i) Amount. Administer 6.8 mL (15 mg altrenogest) per gilt once daily for 14 consecutive days by top-dressing on a portion of each gilt's daily feed.
- (ii) *Indications for use*. For synchronization of estrus in sexually mature gilts that have had at least one estrous cycle.
- (iii) Limitations. Do not use in gilts having a previous or current history of uterine inflammation (i.e., acute, subacute or chronic endometritis). Gilts must not be slaughtered for human consumption for 21 days after the last treatment.

[66 FR 47960, Sept. 17, 2001, as amended at 68 FR 62006, Oct. 31, 2003]

# § 520.62 Aminopentamide hydrogen sulphate tablets.

- (a) Chemical name. 4-(Dimethylamino)-2,2-
- diphenylvaleramide hydrogen sulfate.
  (b) Specifications. Each tablet contains 0.2 milligram of the drug.
- (c) Sponsor. See No. 000856 in §510.600(c) of this chapter.
- (d) Conditions of use. (1) It is intended for use in dogs and cats only for the treatment of vomiting and/or diarrhea, nausea, acute abdominal visceral spasm, pylorospasm, or hypertrophic gastritis.

NOTE: Not for use in animals with glaucoma because of the occurrence of mydriasis.

(2) Dosage is administered by oral tablet every 8 to 12 hours, as follows:

Weight of animal in pounds	Dosage in milligrams
Up to 10	0.1 0.2
21 to 50	0.3
51 to 100	0.4
Over 100	0.5

Dosage may be gradually increased up to a maximum of five times the suggested dosage. Oral administration of tablets may be preceded by subcutaneous or intramuscular use of the injectable form of the drug.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988]

## § 520.82 Aminopropazine fumarate oral dosage forms.

#### § 520.82a Aminopropazine fumarate tablets.

- (a) Specifications. The drug is in tablet form. Each tablet contains aminopropazine fumarate equivalent to 25 milligrams of aminopropazine base.
- (b) *Sponsor*. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use. (1) The drug is used in dogs and cats for reducing excessive smooth muscle contractions, such as occur in urethral spasms associated with urolithiasis.
- (2) It is administered at a dosage level of 1 to 2 milligrams per pound of body weight. The dosage can be repeated every 12 hours, as indicated.<sup>1</sup>
- (3) Not for use in animals intended for food purposes.
- (4) For use only by or on the order of a licensed veterinarian.<sup>1</sup>

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

## § 520.82b Aminopropazine fumarate, neomycin sulfate tablets.

- (a) Specifications. The drug is in tablet form. Each tablet contains both aminopropazine fumarate equivalent to 25 milligrams of aminopropazine base and neomycin sulfate equivalent to 50 milligrams of neomycin base.
- (b) Sponsor. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use. (1) The drug is used in dogs to control bacterial diarrhea caused by organisms susceptible